



## Alka-Seltzer Plus Severe Cold PowerFast Fizz Day & Night Effervescent Tablets Co-Packaged Product

### Alka-Seltzer Plus® Severe Cold PowerFast Fizz Day Effervescent Tablets

#### **Drug Facts**

<b>Active ingredients (in each tablet)</b>	<b>Purposes</b>
Aspirin 325 mg (NSAID)*.....	Pain reliever/fever reducer
Dextromethorphan hydrobromide 10 mg.....	Cough suppressant
Phenylephrine bitartrate 7.8 mg.....	Nasal decongestant

\*nonsteroidal anti-inflammatory drug

#### **Uses**

- temporarily relieves these symptoms due to a cold with cough:
  - minor aches and pains
  - headache
  - cough
  - nasal and sinus congestion
  - sore throat
- temporarily reduces fever

#### **Warnings**

**Reye's syndrome:** Children and teenagers who have or are recovering from chicken pox or flu-like symptoms should not use this product. When using this product, if changes in behavior with nausea and vomiting occur, consult a doctor because these symptoms could be an early sign of Reye's syndrome, a rare but serious illness.

**Allergy alert:** Aspirin may cause a severe allergic reaction which may include:

- hives
- facial swelling
- asthma (wheezing)
- shock

**Stomach bleeding warning:** This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you

- are age 60 or older
- have had stomach ulcers or bleeding problems
- take a blood thinning (anticoagulant) or steroid drug
- take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others)
- have 3 or more alcoholic drinks every day while using this product
- take more or for a longer time than directed

**Sore throat warning:** If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

#### **Do not use**

- if you are allergic to aspirin or any other pain reliever/fever reducer
- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.
- if you have ever had an allergic reaction to this product or any of its ingredients
- in children under 12 years of age

#### **Ask a doctor before use if**

- stomach bleeding warning applies to you
- you have a history of stomach problems, such as heartburn
- you have high blood pressure, heart disease, liver cirrhosis, or kidney disease
- you are taking a diuretic
- you have
  - asthma
  - thyroid disease
  - diabetes
  - cough with excessive phlegm (mucus)
  - difficulty in urination due to enlargement of the prostate gland
  - persistent or chronic cough such as occurs with smoking, asthma, or emphysema
- a sodium-restricted diet



**Alka-Seltzer Plus Severe Cold PowerFast Fizz Day & Night  
Effervescent Tablets  
Co-Packaged Product**

**Ask a doctor or pharmacist before use if you are**

- taking a prescription drug for
  - gout
  - diabetes
  - arthritis

**When using this product do not exceed recommended dosage**

**Stop use and ask a doctor if**

- an allergic reaction occurs. Seek medical help right away.
- you experience any of the following signs of stomach bleeding
  - feel faint
  - vomit blood
  - have bloody or black stools
  - have stomach pain that does not get better
- pain, cough, or nasal congestion gets worse or lasts more than 7 days
- fever gets worse or lasts more than 3 days
- redness or swelling is present
- new symptoms occur
- ringing in the ears or loss of hearing occurs
- cough comes back or occurs with rash or headache that lasts. These could be signs of a serious condition.
- nervousness, dizziness, or sleeplessness occurs

**If pregnant or breast-feeding, ask a health professional before use.**

**It is especially important not to use aspirin during the last 3 months of pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.**

**Keep out of reach of children.** In case of overdose, get medical help or contact a Poison Control Center right away.

**Directions**

- do not take more than the recommended dose
- do not take the Day and Night products at the same time; wait 6 hours after the last Night dose before starting Day product
- adults and children 12 years and over: take 2 tablets fully dissolved in 4 oz of water every 4 hours. Do not exceed 4 tablets in 12 hours or as directed by a doctor.
- children under 12 years: do not use

**Other information**

- **each tablet contains:** potassium 80 mg; sodium 356 mg
- store at room temperature. Avoid excessive heat.

**Inactive ingredients** anhydrous citric acid, calcium silicate, dimethicone, FD&C red #40, FD&C yellow #6, flavor, mannitol, potassium bicarbonate, povidone, sodium bicarbonate, sucralose

**Questions or comments? 1-800-986-0369 (Mon – Fri 9AM – 5PM EST)**



## Alka-Seltzer Plus Severe Cold PowerFast Fizz Day & Night Effervescent Tablets Co-Packaged Product

### Alka-Seltzer Plus® Cold PowerFast Fizz Night Effervescent Tablets

**Drug Facts**

<b>Active ingredients (in each tablet)</b>	<b>Purposes</b>
Aspirin 500 mg (NSAID)*.....	Pain reliever/fever reducer
Dextromethorphan hydrobromide 10 mg.....	Cough suppressant
Doxylamine succinate 6.25 mg.....	Antihistamine
Phenylephrine bitartrate 7.8 mg.....	Nasal decongestant
*nonsteroidal anti-inflammatory drug	

**Uses**

- temporarily relieves these symptoms due to a cold:
  - minor aches and pains
  - headache
  - runny nose
  - sinus congestion and pressure
  - cough
  - sneezing
  - sore throat
  - nasal congestion
- temporarily reduces fever

**Warnings**

**Reye's syndrome:** Children and teenagers who have or are recovering from chicken pox or flu-like symptoms should not use this product. When using this product, if changes in behavior with nausea and vomiting occur, consult a doctor because these symptoms could be an early sign of Reye's syndrome, a rare but serious illness.

**Allergy alert:** Aspirin may cause a severe allergic reaction which may include:

- hives
- facial swelling
- asthma (wheezing)
- shock

**Stomach bleeding warning:** This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you

- are age 60 or older
- have had stomach ulcers or bleeding problems
- take a blood thinning (anticoagulant) or steroid drug
- take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others)
- have 3 or more alcoholic drinks every day while using this product
- take more or for a longer time than directed

**Sore throat warning:** If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

**Do not use to sedate children.**

---

**Do not use**

- if you are allergic to aspirin or any other pain reliever/fever reducer
- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.
- if you have ever had an allergic reaction to this product or any of its ingredients
- in children under 12 years of age

**Ask a doctor before use if**

- stomach bleeding warning applies to you
- you have a history of stomach problems, such as heartburn



## Alka-Seltzer Plus Severe Cold PowerFast Fizz Day & Night Effervescent Tablets Co-Packaged Product

- you have high blood pressure, heart disease, liver cirrhosis, or kidney disease
- you are taking a diuretic
- you have
  - asthma
  - diabetes
  - thyroid disease
  - glaucoma
- cough with excessive phlegm (mucus)
- a breathing problem such as emphysema or chronic bronchitis
- difficulty in urination due to enlargement of the prostate gland
- persistent or chronic cough such as occurs with smoking, asthma, or emphysema
- a sodium-restricted diet

---

**Ask a doctor or pharmacist before use if you are**

- taking a prescription drug for
  - gout
  - diabetes
  - arthritis
- taking sedatives or tranquilizers

---

**When using this product**

- **do not exceed recommended dosage**
- may cause marked drowsiness
- avoid alcoholic drinks
- alcohol, sedatives, and tranquilizers may increase drowsiness
- be careful when driving a motor vehicle or operating machinery
- excitability may occur, especially in children

---

**Stop use and ask a doctor if**

- an allergic reaction occurs. Seek medical help right away.
- you experience any of the following signs of stomach bleeding
  - feel faint
  - vomit blood
  - have bloody or black stools
  - have stomach pain that does not get better
- pain, cough, or nasal congestion gets worse or lasts more than 7 days
- fever gets worse or lasts more than 3 days
- redness or swelling is present
- new symptoms occur
- ringing in the ears or loss of hearing occurs
- cough comes back or occurs with rash or headache that lasts. These could be signs of a serious condition.
- nervousness, dizziness, or sleeplessness occurs

---

**If pregnant or breast-feeding**, ask a health professional before use.

**It is especially important not to use aspirin during the last 3 months of pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.**

**Keep out of reach of children.** In case of overdose, get medical help or contact a Poison Control Center right away.



**Alka-Seltzer Plus Severe Cold PowerFast Fizz Day & Night  
Effervescent Tablets  
Co-Packaged Product**

***Directions***

- do not take more than the recommended dose
- do not take the Day and Night products at the same time; wait 4 hours after the last Day dose before starting Night product
- adults and children 12 years and over: take 2 tablets fully dissolved in 4 oz of water every 6 hours. Do not exceed 4 tablets in 12 hours or as directed by a doctor.
- children under 12 years: do not use

***Other information***

- **each tablet contains:** potassium 80 mg; sodium 356 mg
- store at room temperature. Avoid excessive heat.

***Inactive ingredients*** anhydrous citric acid, calcium silicate, dimethicone, flavors, mannitol, potassium bicarbonate, povidone, sodium bicarbonate, sucralose

***Questions or comments?*** 1-800-986-0369 (Mon-Fri 9AM – 5PM EST)